

PHYS-3301



## Lecture 8

Sep. 17, 2024

## Chapter. 5 Bound States: Simple Case


### Outline:

- The Schrödinger Equation (for interacting particles)
- Stationary States
- Physics Conditions: Well-Behaved Functions
- A Review of Classical Bound States
- Case 1: Particles in a Box – The Infinite Well
- Case 2: The Finite Well
- Case 3: The Simple Harmonic Oscillator
- Expectation Values, Uncertainties, and Operators



Bound states is one in which a particle's motion is restricted by an external force to finite region of space

In Quantum Mechanics –  
Bound States are Standing  
Waves



Not  
forbidden

In Quantum Mechanics –  
Bound States are Standing  
Waves

The  
“Ground state”  
the lowest  
energy state  
is not  
at E=0

Not  
forbidden

In Quantum Mechanics –  
Bound States are Standing  
Waves

The  
“Ground state”  
- the lowest  
energy state  
is not  
at E=0

Consistent with the  
Uncertainty Relations:

$$\Delta x \Delta p_x \geq \frac{\hbar}{2}$$

## The Schrodinger Equation for Interacting Particles

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x, t)}{\partial x^2} = i\hbar \frac{\partial \Psi(x, t)}{\partial t}$$

For free  
particles

or in the absence  
of external forces

$$Ae^{i(kx - \omega t)}$$

$$-\frac{\hbar^2}{2m} (ik)^2 Ae^{i(kx - \omega t)} = i\hbar (-i\omega) Ae^{i(kx - \omega t)}$$

$$\frac{\hbar^2 k^2}{2m} \Psi(x, t) = \hbar \omega \Psi(x, t)$$

$$\frac{p^2}{2m} \Psi(x, t) = E \Psi(x, t) \rightarrow \text{KE } \Psi(x, t) = E \Psi(x, t)$$

Schrödinger eq. is based on E  
accounting - w/o external  
interactions

Try to add potential energy U(x)

## Adding P.E.

$$(\text{KE} + U(x))\Psi(x, t) = E\Psi(x, t)$$

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x, t)}{\partial x^2} + \underline{U(x)\Psi(x, t)} = i\hbar \frac{\partial \Psi(x, t)}{\partial t}$$

→ Time-dependent Schrödinger Eq.

→ To determine the behavior of particle

in (1) CM: solve  $F = m(d^2r/dt^2)$  for  $r$ , given knowledge of Net external  $F$  on particle

in (2) QM: solve the Schrödinger eq. for  $\psi(x,t)$ , given knowledge of P.E.,  $U(x)$

## The Schrodinger Equation for Interacting Particles

and for

## Stationary Potentials

$$U = U(x)$$

$$U \neq U(t)$$

## Key Assumption:

### Factorization of the wave function

$$\Psi(x, t) = \psi(x)\phi(t)$$

Wave function may be express as a product of ...

Standard Math. Technique;  
“Separation of variables”

### Spatial Part

### Temporal Part

**Q:** Why?, **A:** allows us to break a differential eq. with 2 independent variables (x,t) into simpler eqs. For position & time, separately!!

**What happens with the Schrodinger equation?**

$$\Psi(x, t) = \psi(x)\phi(t)$$

$$-\frac{\hbar^2}{2m} \frac{d^2 \psi(x)}{dx^2} + U(x)\psi(x) = E\psi(x)$$

... and factoring out terms constant w.r.t. the partial derivatives ...

$$-\frac{\hbar^2}{2m} \phi(t) \frac{\partial^2 \psi(x)}{\partial x^2} + \underline{U(x)\psi(x)\phi(t)} = i\hbar \psi(x) \frac{\partial \phi(t)}{\partial t}$$

Divide both sides by  $\psi(x)\phi(t)$

$$-\frac{\hbar^2}{2m} \frac{1}{\psi(x)} \frac{\partial^2 \psi(x)}{\partial x^2} + U(x) = i\hbar \frac{1}{\phi(t)} \frac{\partial \phi(t)}{\partial t}$$

Variables are separate now!!

$$-\frac{\hbar^2}{2m} \frac{1}{\psi(x)} \frac{\partial^2 \psi(x)}{\partial x^2} + U(x) = i\hbar \frac{1}{\phi(t)} \frac{\partial \phi(t)}{\partial t}$$

**t and x are independent**

$$-\frac{\hbar^2}{2m} \frac{1}{\psi(x)} \frac{d^2 \psi(x)}{dx^2} + U(x) = i\hbar \frac{1}{\phi(t)} \frac{d\phi(t)}{dt} = C$$

Consider only case in which P.E. is time-independent

Separation Constant

$$i\hbar \frac{1}{\phi(t)} \frac{d\phi(t)}{dt} = C \rightarrow \frac{d\phi(t)}{dt} = -\frac{iC}{\hbar} \phi(t)$$

**The Temporal Part,  $\phi(t)$**

$$\phi(t) = Ae^{-i(C/\hbar)t}$$

$$E = \hbar\omega = C$$

Solution  
(see Appendix K)

$$Ae^{i(kx - \omega t)} \sim Ae^{-i\omega t}, \omega = C/\hbar$$

$$\phi(t) = e^{-i(E/\hbar)t}$$

**Temporal part**

$$\Psi(x, t) = \psi(x)\phi(t)$$

**Total wave function**

$$\Psi(x, t) = \psi(x)e^{-i(E/\hbar)t}$$

$$\phi(t) = e^{-i(E/\hbar)t}$$

**Temporal part**

$$\Psi(x, t) = \psi(x)\phi(t)$$

**Total wave function**

$$\Psi(x, t) = \psi(x)e^{-i(E/\hbar)t}$$

$$\begin{aligned} \Psi^*(x, t)\Psi(x, t) &= [\psi^*(x)e^{+i(E/\hbar)t}][\psi(x)e^{-i(E/\hbar)t}] \\ &= \psi^*(x)\psi(x) \end{aligned}$$

Oops!! Its time dependence disappears!!

**The probability density is time-independent**

**Stationary States**

i.e. the whereabouts of the particle don't change with time in any observable way

$$\phi(t) = e^{-i(E/\hbar)t}$$

**Temporal part**

$$\Psi(x, t) = \psi(x)\phi(t)$$

**Total wave function**

$$\Psi(x, t) = \psi(x)e^{-i(E/\hbar)t}$$

$$\begin{aligned} \Psi^*(x, t)\Psi(x, t) &= [\psi^*(x)e^{+i(E/\hbar)t}][\psi(x)e^{-i(E/\hbar)t}] \\ &= \psi^*(x)\psi(x) \end{aligned}$$

Oops!! Its time dependence disappears!!

**The probability density is time-independent**

**Stationary States**

Quantum Mechanically, electron is not an accelerating charged particles, but rather a stationary "cloud"

## The spatial part of $\psi(x,t)$

Replace C by E, multiply both sides by  $\psi(x)$ ;

## The time-independent Schrodinger equation:

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + U(x)\psi(x) = E\psi(x)$$

Spatial part

**NOTE:**  $\psi(x)$  is *Real*,  
but  $\psi(x,t)$  is *Complex*, because  
 $\phi(t) = e^{-i\omega t}$